



Knowledge grows

# YaraVita<sup>®</sup> FERRICHEL NINETY<sup>™</sup>

A formulated soluble powder iron with nitrogen and phosphorus

Guaranteed Analysis	
Total Nitrogen (N)	12%
Ammoniacal Nitrogen	12%
Available Phosphate (P <sub>2</sub> O <sub>5</sub> )	13%
Iron (Fe)	9%
Derived from Ammonium Phosphate, Ammonium Sulphate, Ferric Citrate	

The information provided is accurate to the best of Yara's knowledge and belief. Any recommendations are meant as a guide and must be adapted to suit local conditions.



## Benefits

- Formulated for safe application at critical growth stages to satisfy crop requirements
- Widely tank mixable with other crop sprays. Visit [www.tankmix.com/yara](http://www.tankmix.com/yara) for details.
- Proven, reliable performance. Tried and tested on a wide range of crops around the world
- High quality, consistent product. Manufactured to ISO 9001 quality assurance standards
- Easy to use liquid formulation. Pours and disperses easily and quickly into the spray tank.
- High nutrient content means lower application rates reducing handling time and waste packaging

# Product Recommendations

## Typical Crop Recommendations\*

- **Apples, Pears:** 1 to 3 lb/acre as soon as there is sufficient leaf cover to intercept the spray, or as soon as first symptoms appear. Repeat as necessary at 10 to 14 day intervals. Do not apply during flowering. Note: Final application to be made at least one month before harvest. Water rate: 50 gallons/acre.
- **Apricots, Cherries, Nectarines, Peaches, Plums:** 1 to 3 lb/acre as soon as there is sufficient leaf cover to intercept the spray, or as soon as first symptoms appear. Repeat as necessary at 10 to 14 day intervals. Do not apply during flowering. Note: Final application to be made at least one month before harvest. Water rate: 50 gallons/acre
- **Citrus:** 1 to 3 lb/acre during Spring and Autumn flushes, at 10 to 14 day intervals as necessary. Do not apply during flowering. Note: Final application to be made at least one month before harvest. Water rate: 50 gallons/acre.
- **Conifers:** 1 to 3 pounds/acre at start of new season leaf production and again in early autumn. Water rate: 50 to 100 gallons/acre.
- **Courgettes, Cucumber, Curcubits (Field Grown):** 1 to 3 lbs/acre from two weeks after transplanting, or as soon as there is sufficient leaf cover to intercept the spray. Repeat as necessary at 10 to 14 day intervals. Do not apply during flowering. Note: Final application to be made at least one month before harvest. Water rate: 20 gallons/acre
- **Vines:** 1 to 3 lb/acre as soon as there is sufficient leaf cover to intercept the spray, or as soon as first symptoms appear. Repeat as necessary at 10 to 14 day intervals. Do not apply during flowering. Note: Final application to be made at least one month before harvest. Water rate: 50 gallons/acre.
- **Turf:** 1 oz./1000 sq.ft. as soon as growth commences in spring. Repeat at 10 to 14 day intervals as necessary. Water rate: 0.5 gallons/1000 sq.ft.
- **Groundnuts:** 1 to 3 lbs/acre as soon as there is sufficient leaf cover to intercept spray or when symptoms are seen. Do not apply during flowering. Water rate: 20 gallons/acre.
- **Lettuce (Field Grown):** 1 to 3 lb/acre from 2 weeks after transplanting or as soon as there is sufficient leaf cover to intercept the spray. Repeat as necessary at 10 to 14 day intervals. Note: Final application to be made at least one month before harvest. Water rate: 20 gallons/acre.
- **Melons (Field Grown):** 1 to 3 lbs/acre from two weeks after transplanting, or as soon as there is sufficient leaf cover to intercept the spray. Repeat as necessary at 10 to 14 day intervals. Do not apply during flowering. Note: Final application to be made at least one month before harvest. Water rate: 20 gallons/acre.
- **Nuts (Deciduous):** 1 to 3 lbs/acre as soon as there is sufficient leaf cover to intercept the spray, or as soon as first symptoms appear. Repeat as necessary at 10 to 14 day intervals. Do not apply during flowering. Note: Final application to be made at least one month before harvest. Water rate: 50 gallons/acre.
- **Onions:** 1 to 3 lb/acre from 2 weeks after transplanting or as soon as there is sufficient leaf cover to intercept the spray. Repeat as necessary at 10 to 14 day intervals. Note: Final application to be made at least one month before harvest. Water rate: 20 gallons/acre.
- **Peppers (Field Grown):** 1 to 3 lb/acre from 2 weeks after transplanting or as soon as there is sufficient leaf cover to intercept the spray. Repeat as necessary at 10 to 14 day intervals. Do not apply during flowering. Note: Final application to be made at least one month before harvest. Water rate: 20 gallons/acre.
- **Soybeans:** 1 to 3 lbs/acre from 2 weeks after emergence, or as soon as there is sufficient leaf cover to intercept the spray. Repeat as necessary at 10 to 14 day intervals. Note: Final application to be made at least one month before harvest. Do not apply during flowering. Water rate: 20 gallons/acre
- **Squash (Field Grown):** 1 to 3 lbs/acre from two weeks after transplanting, or as soon as there is sufficient leaf cover to intercept the spray. Repeat as necessary at 10 to 14 day intervals. Do not apply during flowering. Note: Final application to be made at least one month before harvest. Water rate: 20 gallons/acre.
- **Sugar Cane:** 1 to 3 lb/acre when plants are 12" tall and again 10 to 14 days later. Water rate: 20 gallons/acre.
- **Sweet Corn:** 1 to 3 lb/acre from 2 weeks after transplanting or as soon as there is sufficient leaf cover to intercept the spray. Repeat as necessary at 10 to 14 day intervals. Note: Final application to be made at least one month before harvest. Water rate: 20 gallons/acre.
- **Tomatoes (Field Grown):** 1 to 3 lb/acre from 2 weeks after transplanting or as soon as there is sufficient leaf cover to intercept the spray. Repeat as necessary at 10 to 14 day intervals. Do not apply during flowering. Note: Final application to be made at least one month before harvest. Water rate: 20 gallons/acre.
- **Water Melons (Field Grown):** 1 to 3 lbs/acre from two weeks after transplanting, or as soon as there is sufficient leaf cover to intercept the spray. Repeat as necessary at 10 to 14 day intervals. Do not apply during flowering. Note: Final application to be made at least one month before harvest. Water rate: 20 gallons/acre
- **Zucchini (Field Grown):** 1 to 3 lbs/acre from two weeks after transplanting, or as soon as there is sufficient leaf cover to intercept the spray. Repeat as necessary at 10 to 14 day intervals. Do not apply during flowering. Note: Final application to be made at least one month before harvest. Water rate: 20 gallons/acre

\*The information provided is accurate to the best of Yara's knowledge and belief. Any recommendations are meant as a guide and must be adapted to suit local conditions. Always read the label before use.