



Knowledge grows

YaraVita[®] PHOZTRAC[™]

A formulated product for the treatment of phosphorus, calcium and zinc deficiencies by foliar application

Guaranteed Analysis	
Total Nitrogen (N)	4%
Urea Nitrogen	4%
Available Phosphate (P ₂ O ₅)	15%
Calcium (Ca)	13%
Zinc (Zn)	7.7%
Derived from Urea, Calcium Phosphate, Calcium Carbonate, Zinc Phosphate	

The information provided is accurate to the best of Yara's knowledge and belief. Any recommendations are meant as a guide and must be adapted to suit local conditions.



Benefits

- Formulated for safe application at critical growth stages to satisfy crop requirements
- Widely tank mixable with other crop sprays. Visit www.tankmix.com/yara for details.
- Proven, reliable performance. Tried and tested on a wide range of crops around the world
- High quality, consistent product. Manufactured to ISO 9001 quality assurance standards
- Easy to use liquid formulation. Pours and disperses easily and quickly into the spray tank.
- High nutrient content means lower application rates reducing handling time and waste packaging

Product Recommendations

Typical Crop Recommendations*

- **Alfalfa:** 2 quarts/acre when sufficient cover to intercept spray, and also at flower bud formation. Water rate: 20 gallons/acre.
- **Apples:** 2 to 4 quarts/acre commencing at petal fall with repeat applications at 10 to 14 day intervals. On russet sensitive varieties delay applications until 6 weeks after petal fall. Water rate: 50 gallons/acre.
- **Apricots:** 4 quarts/acre applied 2 to 5 times. First application at pink bud. Thereafter at petal fall with repeat applications at 7 to 14 day intervals. Water rate: 50 to 100 gallons/acre
- **Beans:** 1 to 2 quarts/acre before flowering. Water rate: 25 gallons/acre.
- **Blueberries:** 2 to 4 quarts/acre to new leaf growth in the spring. Water rate: 50 gallons/acre
- **Cole Crops (Broccoli, Brussel Sprouts, Cabbage, Calabrese, Cauliflower):** 2 to 4 quarts/acre applied 2 or 3 times at stem extension/head development with 10 to 14 day intervals. Water rate: 50 gallons per acre.
- **Canola:** 2 to 4 quarts/acre applied 2 to 3 times - at 4 to 6 leaf stage, onset of stem extension, and 10 to 14 days later. Water rate: 25 gallons/acre.
- **Carrots:** 1 to 2 quarts/acre applied 2 or 3 times when crop is 6 inches tall and at 7 to 10 day intervals. Water rate: 25 gallons/acre
- **Celery:** 2 quarts/acre at the 4 to 6 leaf stage. Repeat 10 to 14 days later if necessary. Water rate: 3 to 50 gallons/acre.
- **Cereals:** 2 to 4 quarts/acre at mid-tillering to second node detectable (Zadok's G.S. 25 to 32). Water rate: 25 gallons/acre.
- **Cherries:** 4 quarts/acre applied 2 to 5 times. First application at pink bud. Thereafter at petal fall with repeat applications at 7 to 14 day intervals. Water rate: 50 to 100 gallons/acre
- **Citrus:** 2 to 4 quarts/acre applied 2 to 3 times starting at fruit set with repeat applications at 10 to 14 day intervals. Water rate: 50 to 100 gallons/acre.
- **Conifers:** 1 gallon/acre at the start of new season leaf production, and again in early autumn. Water rate: 3 to 50 gallons/acre.
- **Corn:** 1 to 2 quarts/acre at 4 to 8 leaf stage. Water rate: 25 gallons/acre.
- **Cotton:** 1 to 2 quarts/acre from 10% flowering. Repeat if necessary at 10 to 14 day intervals. Water rate: 30 gallons/acre.
- **Cucurbits (Field Grown):** 1 to 2 quarts/acre at start of flowering. Repeat if necessary at 10 to 14 day intervals. Water rate: 50 gallons/acre.
- **Dry Bean:** 2 quarts per acre applied before flowering. Water rate: 3 to 25 gallons/acre.
- **Garlic:** 2 quarts/acre at the 6 inches stage, and again 10 to 14 days later. Water rate: 3 to 20 gallons/acre.
- **Ginseng:** 1 gallon/acre applied to early season regrowth. Repeat if necessary 10 to 14 days later. Water rate: 50 gallons/acre.
- **Groundnuts:** 2 quarts/acre at the 4 to 6 leaf stage. Consider a second application 10 to 14 days later. Water rate: 20 gallons/acre
- **Lettuce (Field Grown):** 1 to 2 quarts/acre applied 2 or 3 times commencing 10 to 14 days after transplanting or emergence with 7 to 10 day intervals between applications. Water rate: 50 gallons/acre.
- **Nectarines:** 4 quarts/acre applied 2 to 5 times. First application at pink bud. Thereafter at petal fall with repeat applications at 7 to 14 day intervals. Water rate: 50 to 100 gallons/acre.
- **Nursery/Ornamentals:** Two and a half gallons in 100 gallons water (2.5% v/v) as soon as there is sufficient leaf area to intercept a spray. Repeat at 10 to 14 day intervals as necessary. Avoid applications during flowering. Spray a maximum of three applications per crop per annum. Note: Do not apply within one month of picking/marketing. Maximum water rate: 20 gallons/acre.
- **Nuts (Deciduous):** 1 gallon/acre 7 days after petal fall. Repeat if necessary 10 to 14 days later. Water rate: 50 gallons/acre
- **Onions:** 1 to 2 quarts/acre applied once or twice during bulb filling with a 10 to 14 day interval between sprays. Water rate: 25 gallons/acre.
- **Peaches:** 4 quarts/acre applied 2 to 5 times. First application at pink bud. Thereafter at petal fall with repeat applications at 7 to 14 day intervals. Water rate: 50 to 100 gallons/acre
- **Pears:** 4 quarts/acre commencing at petal fall with repeat applications at 10 to 14 day intervals. On russet sensitive varieties delay applications until 6 weeks after petal fall. Water rate: 50 gallons/acre.
- **Peas:** 1 to 2 quarts/acre before flowering. Water rate: 25 gallons/acre.
- **Peppers (Field Grown):** 2 quarts/acre applied 2 to 4 times commencing from flowering on second truss. Repeat at 10 to 14 day intervals. Water rate: 50 gallons/acre.
- **Plums:** 4 quarts/acre applied 2 to 5 times. First application at pink bud. Thereafter at petal fall with repeat applications at 7 to 14 day intervals. Water rate: 50 to 100 gallons/acre
- **Potatoes:** 4 quarts/acre 1 week after 100% crop emergence. Repeat as necessary at 10 to 14 day intervals. Water rate: 25 gallons/acre.
- **Raspberry:** 4 quarts/acre to new leaf growth in the spring. Water rate: 50 gallons/acre.
- **Soybeans:** 2 quarts/acre before flowering. Water rate: 25 gallons/acre.
- **Strawberries (Field Grown):** 4 quarts/acre to new leaf growth in the spring. Water rate: 50 gallons/acre.
- **Sugar Beet:** 2 quarts/acre at 4 to 6 leaf stage. Repeat if necessary at 10 to 14 day intervals. Water rate: 25 gallons/acre.
- **Sugar Cane:** 1 gallon/acre when the plants are 12 inches tall and again 10 to 14 days later. Water rate: 50 gallons/acre
- **Sunflower:** 2 quarts/acre at the 3 to 4 leaf stage. Water rate: 25 gallons/acre
- **Sweet Potatoes:** 2 quarts/acre one week after 100% emergence or transplanting. Repeat applications during tuber bulking at 10 to 14 day intervals. Also, apply at the same rate following recommendation from analysis. Water rate: 20 gallons/acre
- **Tomatoes (Field Grown):** 2 quarts/acre applied 2 to 4 times commencing from flowering on second truss. Repeat at 10 to 14 day intervals. Water rate: 50 gallons/acre.
- **Turf:** 1.5 to 3 fl.oz./1000 sq.ft. as soon as growth commences in spring. Repeat sprays at 10 to 14 day intervals as necessary. Water rate: 0.5 gallons/1000sq.ft.
- **Vines:** 2 to 4 quarts/acre when new shoots are at the 4 to 6 leaf stage. Repeat if necessary at 10 to 14 day intervals. Water rate: 50 gallons/acre.

*The information provided is accurate to the best of Yara's knowledge and belief. Any recommendations are meant as a guide and must be adapted to suit local conditions. Always read the label before use.